arXiv:1607.03322v1 [math.RA] 12 Jul 2016

A NATURAL MAP FROM A QUANTIZED SPACE ONTO ITS SEMICLASSICAL LIMIT

SEI-QWON OH

Abstract. Let $\widehat{\Gamma}$ be the natural map given in [11, §1]. Here, we construct a deformation B_q of a Poisson algebra B_1 and a prime ideal P of B_q such that $\widehat{\Gamma}(P)$ is not a Poisson prime ideal of B_1.

Recall the star product in [8, 1.1]. Let $\hbar$ be a formal variable. Let $\mathbf{k}$ be a field of characteristic zero and let A be a Poisson algebra over $\mathbf{k}$. The star product $\star$ on $A[[\hbar]]$ is given as follows: for any $f = \sum_{n\geq 0} f_n\hbar^n, g = \sum_{n\geq 0} g_n\hbar^n \in A[[\hbar]]$,

$$f \star g := \sum_{k,l\geq 0} f_k g_l \hbar^{k+l} + \sum_{k,l\geq 0, m\geq 1} B_m(f_k, g_l)\hbar^{k+l+m}$$

such that

$$\frac{f\star g - g\star f}{\hbar}\,|_{\hbar=0} = \{f_0, g_0\},$$

where $B_i : A \times A \longrightarrow A$ are differential operators with respect to each argument. The $\mathbf{k}[[\hbar]]$-algebra $A[[\hbar]]$ with a star product is called a *quantization* of A.

Let B be a $\mathbf{k}[\hbar]$-subalgebra of $A[[\hbar]]$ containing A. Note that $\hbar$ is a nonzero, nonunit and non-zero-divisor such that $B/\hbar B \cong A$ is commutative. For $\lambda \in \mathbf{k}$, $(\hbar - \lambda)B$ is an ideal of B and thus $B_\lambda := B/(\hbar-\lambda)B$ is a $\mathbf{k}$-algebra. In particular, we can recover the Poisson algebra $B_0 \cong A$ and $B_\lambda = 0$ if $\hbar - \lambda$ is a unit in B. Suppose that $0 \neq \lambda \in \mathbf{k}$ such that $\hbar - \lambda$ is a nonunit in B. Then B_λ is a non-trivial $\mathbf{k}$-algebra. (Let $\lambda \in \mathbf{k}$ be nonzero. Then $\hbar - \lambda$ is a unit in $A[[\hbar]]$. Thus, if $\hbar - \lambda$ is a nonunit in B, then B is a proper subalgebra of $A[[\hbar]]$.)

Let t be an indeterminate. In the above paragraphes, replace $\mathbf{k}[[h]]$ by $\mathbf{k}[t, t^{-1}]$ and let B be any $\mathbf{k}[t, t^{-1}]$-algebra. Assume that $t-1$ is a nonzero, nonunit and non-zero-divisor in B such that $B_1 := B/(t-1)B$ is commutative as $\hbar$ is such an element in $A[[\hbar]]$. It is easy to check that B_1 is a Poisson algebra with Poisson bracket

$$\{\overline{a}, \overline{b}\} = \overline{(t-1)^{-1}(ab-ba)} \tag{1}$$

for all $\overline{a} = a+(t-1)B, \overline{b} = b+(t-1)B \in B_1$. The Poisson algebra B_1 plays a role as the Poisson algebra A in the first paragraph. The Poisson algebra B_1 is called a *semiclassical limit* of B. For $\lambda \in \mathbf{k}$, set $B_\lambda := B/(t-\lambda)B$ and, for any $\mathbf{k}$-algebra R, denote by $\mathrm{GKdim}(R)$ the Gelfand-Kirillov dimension of R. (Refer to [9] for the Gelfand-Kirillov dimension.) The $\mathbf{k}$-algebra B_λ is called a *deformation* of B_1. Let $\mathbb{K}$ be a nonempty subset of the set

$$\{1, 0 \neq \lambda \in \mathbf{k} \mid t-\lambda \text{ is a nonunit in } B \text{ and } \mathrm{GKdim}(B_\lambda)=\mathrm{GKdim}(B_1)\}.$$

2000 *Mathematics Subject Classification.* 17B63, 16S36.

Key words and phrases. Poisson algebra, quantization, deformation, semiclassical limit,

That is, if $\lambda \in \mathbb{K}$ then $B_\lambda \neq 0$ and $\mathrm{GKdim}(B_\lambda)$ is equal to $\mathrm{GKdim}(B_1)$. This article concerns with the natural map from a family of deformations of B_1, $\{B_\lambda | \lambda \in \mathbb{K}\}$, into the Poisson algebra B_1 given in [11, §1]. Recall the natural map given in [11, §1]. Define a natural homomorphism of $\mathbf{k}$-algebras

$$\gamma : B \longrightarrow \prod_{\lambda \in \mathbb{K}} B_\lambda, \quad b \mapsto \gamma(b) := (b + (t-\lambda)B)_{\lambda \in \mathbb{K}}.$$

Let $\gamma_1 : B \longrightarrow B_1$ be the natural projection given by $\gamma_1(b) = b + (t-1)B$ for all $b \in B$. If γ is a monomorphism, then there exists the composition

$$\Gamma = \gamma_1 \circ \gamma^{-1} : \gamma(B) \longrightarrow B_1.$$

Note that Γ is surjective. Denote by q a parameter taking values in $\mathbb{K}$. Then the $\mathbf{k}$-algebra B_q presents a family of deformations of B_1, $\{B_\lambda | \lambda \in \mathbb{K}\}$, and there is a natural map

$$\widehat{\ } : B_q \longrightarrow \prod_{\lambda \in \mathbb{K}} B_\lambda, \quad z(q) \mapsto (z(\lambda))_{\lambda \in \mathbb{K}}$$

such that the image $\widehat{q}$ of q by the map $\widehat{\ }$ is equal to $\gamma(t)$. Note that the image $\widehat{B_q}$ of B_q by the map $\widehat{\ }$ is contained in $\gamma(B)$. Hence there is a composition

$$\widehat{\Gamma} = \widehat{\ } \circ \Gamma : B_q \longrightarrow B_1,$$

which is the natural map given in [11, §1].

Deformations B_q of B_1 share algebraic properties with the Poisson algebra B_1 in many cases, for instance, prime ideals and Poisson prime ideals, primitive ideals and Poisson primitive ideals, central elements and Poisson central elements, automorphisms and Poisson automorphisms. (See [1], [3], [5], [7], [10].) Let I be an ideal of B_q. It is proved in [11, Theorem 1.4] that the image $\widehat{\Gamma}(I)$ of I by the map $\widehat{\Gamma}$ is a Poisson ideal of B_1. Moreover, it is shown in [11] under certain conditions that the natural map $\widehat{\Gamma}$ induces a homeomorphism between the spectrum of B_q and the Poisson spectrum of B_1 for the case of the quantized affine algebra and the quantized Weyl algebra.

The aim of this article is to give an example such that the natural map $\widehat{\Gamma}$ does not preserves primeness. That is, we construct deformations B_q of a Poisson algebra B_1 and a prime ideal P of B_q such that $\widehat{\Gamma}(P)$ is not a Poisson prime ideal of B_1.

Assume throughout the article that $\mathbf{k}$ denotes a field of characteristic zero and that all algebras have unity. A commutative algebra A is said to be a *Poisson algebra* if there exists a bilinear product $\{-,-\}$ on A, called a *Poisson bracket*, such that $(A, \{-,-\})$ is a Lie algebra and $\{ab, c\} = a\{b,c\} + \{a,c\}b$ for all $a, b, c \in A$. An ideal I of a Poisson algebra A is said to be a *Poisson ideal* if $\{I, A\} \subseteq I$. A Poisson ideal P is said to be *Poisson prime* if, for all Poisson ideals I and J, $IJ \subseteq P$ implies $I \subseteq P$ or $J \subseteq P$. If A is noetherian then a Poisson prime ideal of A is a prime ideal by [4, Lemma 1.1(d)].

Let B be the $\mathbf{k}[t, t^{-1}]$-algebra generated by e, f, h with the relations

$$ef - fe = (t-1)h, \quad he - eh = 2(t-1)e, \quad hf - fh = -2(t-1)f.$$

Note that B is an iterated skew polynomial ring

$$B = \mathbf{k}[t, t^{-1}][h][e; \alpha_1][f; \alpha_2, \delta_2]$$

for suitable maps $\alpha_1, \alpha_2, \delta_2$. (Refer to [6, Chapter 2] for the skew polynomial ring.) Hence $t-1$ is a non-zero-divisor in B. Clearly, $t-1$ is a nonzero and nonunit such that $B_1 := B/(t-1)B$ is commutative and thus the semiclassical limit B_1 of B is the Poisson algebra $\mathbf{k}[e, f, h]$ with Poisson bracket

$$\{e, f\} = h, \quad \{h, e\} = 2e, \quad \{h, f\} = -2f \tag{2}$$

by (1). Set

$$\mathbb{K} := \mathbf{k} \setminus (\{0, 1\} \cup \{\text{roots of unity}\}).$$

Then $\mathbb{K}$ is an infinite set clearly and $t - \lambda$, $\lambda \in \mathbb{K}$, is a nonunit in B. Denote by B_λ the deformation $B/(t-\lambda)B$ of B_1. Note that, for each $\lambda \in \mathbb{K}$, B_λ is the $\mathbf{k}$-algebra generated by e, f, h with the relations

$$ef - fe = (\lambda - 1)h, \quad he - eh = 2(\lambda - 1)e, \quad hf - fh = -2(\lambda - 1)f \tag{3}$$

and observe that B_λ is an iterated skew polynomial ring

$$B_\lambda = \mathbf{k}[h][e; \alpha_1'][f; \alpha_2', \delta_2']$$

for suitable maps $\alpha_1', \alpha_2', \delta_2'$. Hence the set of standard monomials

$$\{e^i f^j h^k | i, j, k = 0, 1, \cdots\}$$

forms a $\mathbf{k}[t, t^{-1}]$-basis of B and a $\mathbf{k}$-basis of B_λ for each $\lambda \in \mathbf{K}$. It is also a $\mathbf{k}$-basis of B_1. Hence GKdim(B_λ)=GKdim(B_1)= 3 for each $\lambda \in \mathbb{K}$ by [9, Example 3.6]. By [11, Lemma 1.2], there exists a monomorphism of $\mathbf{k}$-algebras

$$\gamma : B \longrightarrow \prod_{\lambda \in \mathbb{K}} B_\lambda$$

and thus there exists the composition

$$\Gamma = \gamma_1 \circ \gamma^{-1} : \gamma(B) \longrightarrow B_1,$$

where γ_1 is the canonical projection

$$\gamma_1 : B \longrightarrow B_1, \quad b \mapsto b + (t-1)B.$$

Note that Γ is an epimorphism of $\mathbf{k}$-algebras.

Let q be not a root of unity. Then B_q is the $\mathbf{k}$-algebra generated by e, f, h with the relations

$$ef - fe = (q-1)h, \quad he - eh = 2(q-1)e, \quad hf - fh = -2(q-1)f, \tag{4}$$

which is obtained from (3) by replacing λ by q. Moreover, q plays a role as a parameter taking values in $\mathbb{K}$ and thus B_q presents a family of deformations of B_1, $\{B_\lambda | \lambda \in \mathbb{K}\}$. There exists a map

$$\widehat{\ } : B_q \longrightarrow \prod_{\lambda \in \mathbb{K}} B_\lambda, \quad z(q) \mapsto (z(\lambda))_{\lambda \in \mathbb{K}}.$$

Observe that the image $\widehat{B_q}$ of B_q by $\widehat{\ }$ is contained in $\gamma(B)$. Thus there exists the composition

$$\widehat{\Gamma} = \widehat{\ } \circ \Gamma : B_q \longrightarrow B_1.$$

Note that $\widehat{\Gamma}$ is a surjective map but not a $\mathbf{k}$-linear map since $\widehat{\Gamma}(q) = 1$.

The universal enveloping algebra $U(\mathfrak{sl}_2(\mathbf{k}))$ of the Lie algebra $\mathfrak{sl}_2(\mathbf{k})$ is the $\mathbf{k}$-algebra generated by E, F, H subject to the relations

$$EF - FE = H, \quad HE - EH = 2E, \quad HF - FH = -2F. \tag{5}$$

One should compare (2) with (5). The map from $U(\mathfrak{sl}_2(\mathbf{k}))$ onto B_q defined by

$$E \mapsto (q-1)^{-1}e, \quad F \mapsto (q-1)^{-1}f, \quad H \mapsto (q-1)^{-1}h \tag{6}$$

is an isomorphism of algebras by (4) and (5). Hence $U(\mathfrak{sl}_2(\mathbf{k})) \cong B_q$.

Note that the element $\Omega = 4ef + h^2 - 2(q-1)h$ of B_q is a central element of B_q which corresponds to a scalar multiple of the Casimir element of $U(\mathfrak{sl}_2(\mathbf{k}))$. For each integer $n \geq 2$, the ideal

$$P_n = \langle e^n, \Omega - (q-1)^2(n^2-1) \rangle$$

of B_q is a prime ideal by (6) and [2, Proposition 4.3]. But $\widehat{\Gamma}(P_n)$ is the Poisson ideal

$$\langle e^n, 4ef + h^2 \rangle$$

of B_1 by [11, Theorem 1.4], which is not prime since $n \geq 2$.

Acknowledgments The author is supported by National Research Foundation of Korea, Grant 2012-007347.

References

1. Alexei Belov-Kanel and Maxim Kontsevich, *Automorphisms of the Weyl algebra*, Lett. Math. Phys. **74** (2005), 181–199.
2. Stefan Catoiu, *Ideals of the enveloping algebra $U(\mathfrak{sl}_2)$*, J. Algebra **202** (1998), 142–177.
3. Eun-Hee Cho and Sei-Qwon Oh, *Semiclassical limits of Ore extensions and a Poisson generalized Weyl algebra*, Lett. Math. Phys. **106** (2016), no. 7, 997–1009.
4. K. R. Goodearl, *A Dixmier-Moeglin equivalence for Poisson algebras with torus actions*, in Algebra and Its Applications (D. V. Huynh, S. K. Jain, and S. R. Lopez-Permouth, Eds.) Contemp. Math. **419** (2006), 131–154.
5. ———, *Semiclassical limits of quantized coordinate rings*, in Advances in Ring Theory (D. V. Huynh and S. R. Lopez-Permouth, Eds.) Basel Birkhäuser (2009), 165–204.
6. K. R. Goodearl and R. B. Warfield, *An introduction to noncommutative noetherian rings*, Second ed., London Mathematical Society Student Text 61, Cambridge University Press, 2004.
7. David A. Jordan and Sei-Qwon Oh, *Poisson spectra in polynomial algebras*, Journal of Algebra **400** (2014), 56–71.
8. M. Kontsevich, *Deformation quantization of Poisson manifolds*, arXiv:q-alg/9709040 (1997). (The final version is available at the website: http://www.ihes.fr/ maxim/TEXTS/DefQuant_final.pdf)
9. G. R. Krause and T. H. Lenagan, *Growth of algebras and Gelfand-Kirillov dimension*, Revised Edition, Graduate Studies in Mathematics, vol. 22, American Mathematical Society, Providence, 1999.
10. Sei-Qwon Oh, *Quantum and Poisson structures of multi-parameter symplectic and Euclidean spaces*, J. Algebra **319** (2008), 4485–4535.
11. Sei-Qwon Oh, *A natural map from a quantized space onto its semiclassical limit and a multi-parameter Poisson Weyl algebra*, Comm. Algebra **45** (2017), DOI 10.1080/00927872.2016.1175448. (available at arXiv:1511.00802v1)

Department of Mathematics, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon 34134, Korea

E-mail address: sqoh@cnu.ac.kr